USDA United States Department of Agriculture

Natural Resources Conservation Service

Montana Water Supply Outlook Report June 1, 2011



Picture: Madison Plateau SNOTEL Site near West Yellowstone

Water Supply Outlook Report
and
Federal - State - Private
Cooperative Snow Surveys

For more water supply and resource management information, contact:

Brian Domonkos
Water Supply Specialist
Federal Building
10 East Babcock, Room 443
Bozeman, MT 59715
Phone 406-587-6991

How forecasts are made

Most of the annual streamflow in the Western United States originates as snowfall that has accumulated high in the mountains during winter and early spring. As the snowpack accumulates, hydrologists estimate the runoff that will occur when it melts. Predictions are based on careful measurements of snow water equivalent at selected index points. Precipitation, temperature, soil moisture and antecedent streamflow data are combined with snowpack data to prepare runoff forecasts. Streamflow forecasts are coordinated by Natural Resources Conservation Service and National Weather Service hydrologists. This report presents a comprehensive picture of water supply conditions for areas dependent upon surface runoff. It includes selected streamflow forecasts, summarized snowpack and precipitation data, reservoir storage data, and narratives describing current conditions.

Snowpack data are obtained by using a combination of manual and automated SNOTEL measurement methods. Manual readings of snow depth and water equivalent are taken at locations called snow courses on a monthly or semi-monthly schedule during the winter. In addition, snow water equivalent, precipitation and temperature are monitored on a daily basis and transmitted via meteor burst telemetry to central data collection facilities. Both monthly and daily data are used to project snowmelt runoff.

Forecast uncertainty originates from two sources: (1) uncertainty of future hydrologic and climatic conditions, and (2) error in the forecasting procedure. To express the uncertainty in the most probable forecast, four additional forecasts are provided. The actual streamflow can be expected to exceed the most probable forecast 50% of the time. Similarly, the actual streamflow volume can be expected to exceed the 90% forecast volume 90% of the time. The same is true for the 70%, 30%, and 10% forecasts. Generally, the 90% and 70% forecasts reflect drier than normal hydrologic and climatic conditions; the 30% and 10% forecasts reflect wetter than normal conditions. As the forecast season progresses, a greater portion of the future hydrologic and climatic uncertainty will become known and the additional forecasts will move closer to the most probable forecast.

The United States Department of Agriculture (USDA) prohibits discrimination in its programs on the basis of race, color, national origin, sex, religion, age, disability, political beliefs and marital or familial status. (Not all prohibited bases apply to all programs.) Persons with disabilities who require alternative means for communication of program information (Braille, large print, audiotape, etc.) should contact USDA's TARGET Center at (202) 720-2600 (voice and TDD).

To file a complaint, write the Secretary of Agriculture, U.S. Department of Agriculture, Washington, D.C., 20250, or call 1-800-245-6340 (voice) or (202) 720-1127 (TDD). USDA is an equal employment opportunity employer.

Mountain Snow Water Equivalent



Surface Water Supply Index (SWSI) Values



RIVER INDEX & SWSI VALUES

1 Marias River above Tiber Reservoir 3.4
2 Tobacco River 3.4
3 Kootenai Ft. Steele to Libby Dam 2.4
4 Kootenai River below Libby Dam
5 Fisher River 3.8
6 Yaak River 3.2
7 North Fk. Flathead River 3.8
8 Middle Fk. Flathead River 3.8
9 South Fk. Flathead River 3.8
10 Flathead River at Columbia Falls 3.2
12 Swan River 3.9
13 Flathead River at Polson 3.8
14 Mission Valley
15 Little Bitterroot River
16 Clark Fork River above Milltown 3.5
17 Blackfoot River 4
18 Clark Fork River above Missoula 3.6
19 Bitterroot River 2.5
20 Clark Fork River below Bitterroot River 3.2
21 Clark Fork River below Flathead River 3.6
22 Beaverhead River 2.3
23 Ruby River 2.4
24 Big Hole River 3.7
25 Boulder River (Jefferson) 3
26 Jefferson River 3
27 Madison River 2.8
28 Gallatin River 3.4
29 Missouri River above Canyon Ferry 3.3
30 Missouri River below Canyon Ferry 3.1
31 Smith River
32 Sun River 3.2
33 Teton River 2.3
34 Birch/Dupuyer Creeks 3.3
35 Marias River 3.9
36 Musselshell River 3.8
37 Missouri River above Fort Peck 2.3
38 Missouri River below Fort Peck 3.9
39 Milk River 3.89
40 Dearborn River near Craig 2.8
41 Yellowstone River above Livingston 3.7
42 Shields River 3.8
43 Boulder River (Yellowstone) 3.6
44 Stillwater River 3.8
45 Rock/Red Lodge Creeks 3.8
46 Clarks Fork River 3.9
47 Yellowstone River above Bighorn River 3.7
48 Bighorn River below Bighorn Lake 4
49 Little Bighorn River 3.8
50 Yellowstone River below Bighorn River 3.8
51 Tongue River 3.9
52 Powder River 4
53 Upper Judith River
54 St. Mary River 3.3

Extremely Dry -4.0 to -3.0
Moderately Dry -2.9 to -2.0
Slightly Dry -1.9 to -1.0
Near Average -0.9 to 0.9
Slightly Wet 1.0 to 1.9
Moderately Wet 2.0 to 2.9
Extremely Wet 3.0 to 4.0
SWSI Not Applicable

June 1, 2011

NOTE: Data used to generate this map are PROVISIONAL and SUBJECT TO CHANGE.

NRCS Natural Resources Conservation Service

SNOTEL AND SNOW COURSE DATA SUMMARY

JUNE 2011

SNOW COURSE	ELEVATION	DATE	SNOW DEPTH	WATER CONTENT	LAST YEAR	AVERAGE 71-00
ALBRO LAKE SNOTEL	8300	6/01/11	80	28.4	22.1	13.4
BADGER PASS SNOTEL	6900	6/01/11	101	49.3	26.2	22.9
BANFIELD MTN SNOTEL	5600	6/01/11	36	15.1	.0	2.9
BARKER LAKES SNOTEL	8250	6/01/11	77	25.6	20.0	9.5
BASIN CREEK SNOTEL	7180	6/01/11	26	8.1	3.4	4.1
BEAGLE SPGS SNOTEL	8850	6/01/11	27	9.3	6.4	1.4
BEAVER CREEK SNOTEL	7850	6/01/11	49	21.2	9.3	8.6
BISSON CREEK SNOTEL	4920	6/01/11	0	.0	.0	.0
BLACK BEAR SNOTEL	7950	6/01/11	114	53.9	22.8	26.1
BLACK PINE SNOTEL	7100	6/01/11	22	8.1	.0	1.9
BLACKTAIL MTN SNOTEL	5650	6/01/11	20	9.2	--	--
BLOODY DICK SNOTEL	7550	6/01/11	24	8.9	.4	1.0
BOULDER MTN SNOTEL	7950	6/01/11	69	27.2	18.0	9.2
BOX CANYON SNOTEL	6700	6/01/11	0	.0	.0	.0
BOXELDER CREEK	5100	6/01/11	0	.0	.0	.2
BRACKETT CR SNOTEL	7320	6/01/11	61	28.1	16.9	5.1
BURNT MTN SNOTEL	5880	6/01/11	0	.0	.0	.0
CALVERT CR SNOTEL	6430	6/01/11	0	.0	.0	.0
CARROT BASIN SNOTEL	9000	6/01/11	90	35.7	22.3	24.2
CHICKEN CREEK	4060	5/26/11	0	.0	.0	.0
CLOVER MDW SNOTEL	8800	6/01/11	59	22.9	18.0	11.3
COLE CREEK SNOTEL	7850	6/01/11	72	25.2	17.4	11.0
COMBINATION SNOTEL	5600	6/01/11	0	.0	.0	.0
COPPER BOTTOM SNOTEL	5200	6/01/11	0	.0	.0	.0
COPPER CAMP SNOTEL	6950	6/01/11	99	48.7	14.0	--
COYOTE HILL	4200	5/27/11	0	.0	.0	--
CRYSTAL LAKE SNOTEL	6050	6/01/11	44	15.6	20.0	.7
DAISY PEAK SNOTEL	7600	6/01/11	42	14.7	5.3	3.4
DALY CREEK SNOTEL	5780	6/01/11	0	.0	.0	.0
DARKHORSE LK. SNOTEL	8700	6/01/11	110	46.4	29.0	26.3
DEADMAN CR SNOTEL	6450	6/01/11	8	3.0	.0	.0
DISCOVERY BASIN	7050	5/27/11	30	10.7	5.7	2.4
DIVIDE SNOTEL	7800	6/01/11	19	6.5	2.8	2.7
DIX HILL	6400	5/29/11	0	.0	.0	--
DUPUYER CREEK SNOTEL	5750	6/01/11	21	5.9	.0	.4
EMERY CREEK SNOTEL	4350	6/01/11	0	.0	.0	.0
FISHER CREEK SNOTEL	9100	6/01/11	120	53.9	27.9	30.1
FLATTOP MTN SNOTEL	6300	6/01/11	142	62.0	38.5	36.5
FROHNER MDWS SNOTEL	6480	6/01/11	2	.7	.0	.7
GARVER CREEK SNOTEL	4250	6/01/11	0	.0	.0	.0
GRAVE CRK SNOTEL	4300	6/01/11	21	8.7	.0	.0
HAND CREEK SNOTEL	5030	6/01/11	0	.0	.0	.0
HAWKINS LAKE SNOTEL	6450	6/01/11	80	40.4	13.9	17.8
HELL ROARING DIVIDE	5770	5/27/11	98	41.4	16.3	10.8
HERRIG JUNCTION	4850	5/26/11	58	26.8	8.7	5.4
HOODOO BASIN SNOTEL	6050	6/01/11	117	53.8	19.6	28.4
KRAFT CREEK SNOTEL	4750	6/01/11	0	.0	.0	.0
LAKEVIEW RDG. SNOTEL	7400	6/01/11	14	4.1	.0	.3
LEMHI RIDGE SNOTEL	8100	6/01/11	40	13.0	9.0	1.4
LICK CREEK SNOTEL	6860	6/01/11	27	10.0	1.4	.5
LONE MOUNTAIN SNOTEL	8880	6/01/11	72	30.8	14.8	11.8
LOWER TWIN SNOTEL	7900	6/01/11	68	23.4	19.8	13.0
LUBRECHT SNOTEL	4680	6/01/11	0	.0	.0	.0
MADISON PLT SNOTEL	7750	6/01/11	72	32.0	9.8	9.0
MANY GLACIER SNOTEL	4900	6/01/11	0	.0	.0	.0
MARIAS PASS	5250	5/30/11	34	15.2	--	--
MONUMENT PK SNOTEL	8850	6/01/11	73	30.3	15.8	15.9
MOSS PEAK SNOTEL	6780	6/01/11	134	63.7	41.8	31.0

SNOW COURSE	ELEVATION	DATE	SNOW DEPTH	WATER CONTENT	LAST YEAR	AVERAGE 71-00
MT LOCKHART SNOTEL	6400	6/01/11	59	27.6	7.9	9.6
MULE CREEK SNOTEL	8300	6/01/11	55	19.4	16.8	9.3
N.E. ENTRANCE SNOTEL	7350	6/01/11	0	.0	.0	.4
NEVADA RIDGE SNOTEL	7020	6/01/11	43	19.2	3.8	3.4
NEZ PERCE CMP SNOTEL	5650	6/01/11	8	4.3	.0	.3
N.F. ELK CR SNOTEL	6250	6/01/11	8	3.1	.0	.6
NF JOCKO SNOTEL	6330	6/01/11	109	53.5	30.7	26.4
NOISY BASIN SNOTEL	6040	6/01/11	140	69.7	38.2	30.1
OPHIR PARK	7150	5/29/11	33	13.9	4.9	--
PETERSON MDW SNOTEL	7200	6/01/11	36	11.4	10.6	2.7
PICKFOOT CRK SNOTEL	6650	6/01/11	12	4.2	.0	.0
PIKE CREEK SNOTEL	5930	6/01/11	29	13.5	.0	7.3
PLACER BASIN SNOTEL	8830	6/01/11	84	34.1	16.1	14.6
POORMAN CR SNOTEL	5100	6/01/11	88	44.5	6.2	11.1
PORCUPINE SNOTEL	6500	6/01/11	0	.0	.0	.0
RED MOUNTAIN	6000	5/31/11	41	19.7	5.9	4.1
REVAIS CREEK	4800	6/01/11	0	.0	.0	.0
ROCKER PEAK SNOTEL	8000	6/01/11	59	23.1	15.0	11.7
ROCKY BOY SNOTEL	4700	6/01/11	0	.0	.0	.2
SACAJAWEA SNOTEL	6550	6/01/11	22	9.7	2.1	.0
SADDLE MTN SNOTEL	7900	6/01/11	73	33.2	12.6	16.3
S.F. SHIELDS SNOTEL	8100	6/01/11	72	27.2	17.3	12.3
SHORT CREEK SNOTEL	7000	6/01/11	0	.0	.0	.0
SHOWER FALLS SNOTEL	8100	6/01/11	89	37.0	25.4	20.5
SKALKAHO SNOTEL	7260	6/01/11	54	25.1	6.9	14.6
SLEEPING WOMAN SNTL	6150	6/01/11	28	12.1	.0	1.2
SPOTTED BEAR MTN.	7000	5/31/11	15	6.1	--	--
SPUR PARK SNOTEL	8100	6/01/11	81	36.8	21.2	14.2
STAHL PEAK SNOTEL	6030	6/01/11	134	57.4	34.8	28.0
STRYKER BASIN	6180	5/26/11	99	46.8	26.9	19.4
STUART MOUNTAIN SNTL	7400	6/01/11	92	46.2	21.2	17.0
TAYLOR ROAD	4080	6/01/11	0	.0	.0	.2
TEPEE CREEK SNOTEL	8000	6/01/11	21	8.6	1.5	5.2
TIZER BASIN SNOTEL	6840	6/01/11	19	5.8	.0	.3
TV MOUNTAIN	6800	5/31/11	40	18.8	6.9	6.8
TWELVEMILE SNOTEL	5600	6/01/11	0	.0	.0	.4
TWENTY-ONE MILE	7150	5/30/11	24	10.7	--	--
TWIN LAKES SNOTEL	6400	6/01/11	75	38.8	12.4	22.3
WALDRON SNOTEL	5600	6/01/11	14	5.3	.0	.1
WARM SPRINGS SNOTEL	7800	6/01/11	77	32.5	24.1	17.0
WEST YELL'ST SNOTEL	6700	6/01/11	0	.0	.0	.0
WHISKEY CREEK SNOTEL	6800	6/01/11	35	13.4	.0	1.2
WHITE MILL SNOTEL	8700	6/01/11	86	35.1	16.2	18.0
WOOD CREEK SNOTEL	5960	6/01/11	27	9.6	.0	.5

Montana Water Supply Outlook Report as of June 1, 2011

With the amount of snow still present in the mountains, and the recent weather patterns, the fact that summer is merely three weeks away is difficult to grasp. La Nina has continued to squeeze its tight grip since November 2010. This has allowed all components that factor into water supply to indicate a well above average outlook for the remainder of this water year. Even reservoir numbers, although below normal levels, reflect well above average conditions. Reservoir levels are currently low in anticipation of the above average snowpack yet to runoff.

Snowpack

The St. Mary and Milk River drainages are the only major basins in the state to be below 200 percent of average snowpack on June 1. All other basins are near 250 percent of average snowpack with some approaching 400 percent of average. The current La Nina weather patterns drove late season snow storms at the end of May, dropping snow at the higher elevations, resulting in new snowpack maximums for the year and period of record. Keep in mind that many snowpacks have typically melted off at this point in time. This coupled with snow still present at many observation sites, results in elevated averages at many locations.

RIVER BASIN	% OF AVERAGE	% OF LAST YEAR	MAY % CHANGE
COLUMBIA	237	235	+77
KOOTENAI, MONTANA	249	289	+80
FLATHEAD, MONTANA	235	196	+55
UPPER CLARK FORK	240	205	+86
BITTERROOT	207	381	+57
LOWER CLARK FORK	288	413	+121
MISSOURI	261	192	+108
MISSOURI HEADWATERS	240	188	+97
JEFFERSON	228	170	+91
MADISON	218	193	+73
GALLATIN	244	187	+101
MISSOURI MAINSTEM	328	195	+148
HEADWATERS MAINSTEM	300	207	+143
SMITH-JUDITH-MUSSELSHELL	386	152	+198
SUN-TETON-MARIAS	273	326	+107
BEARPAW MOUNTAINS	--	--	--
ST. MARY	170	161	+28
ST. MARY & MILK	167	161	+22
YELLOWSTONE	268	219	+123
UPPER YELLOWSTONE	222	217	+71
LOWER YELLOWSTONE	305	227	+166
STATE-WIDE	243	219	+83

Precipitation

May mountain and valley precipitation across the state was 167 percent of average and 125 percent of last year, while the water year precipitation was 131 percent of average and 142 percent of last year. West of the Continental Divide, May mountain and valley precipitation was 114 percent of average and 91 percent of last year and the water year precipitation was 126 percent of average and 152 percent of last year. East of the Divide, May mountain and valley precipitation was 190 percent of average and 138 percent of last year and the water year precipitation was 135 percent of average and 136 percent of last year.

RIVER BASIN	MAY % OF AVERAGE	WATER YEAR % OF AVERAGE
COLUMBIA	114	126
KOOTENAI	115	128
FLATHEAD	109	132
UPPER CLARK FORK	120	124
BITTERROOT	122	119
LOWER CLARK FORK	107	126
MISSOURI	169	132
JEFFERSON	145	121
MADISON	153	123
GALLATIN	113	120
MISSOURI MAINSTEM	196	145
SMITH-JUDITH-MUSSELSHELL	223	153
SUN-TETON-MARIAS	135	124
MILK	173	149
ST. MARY	104	117
YELLOWSTONE	220	137
UPPER YELLOWSTONE	183	134
LOWER YELLOWSTONE	261	146
STATEWIDE	167	131

Reservoirs

State-wide reservoir storage was 105 percent of average and 106 percent of last year. Reservoir storage west of the divide was 79 percent of average and 69 percent of last year. East of the Divide, reservoir storage was 115 percent of average and 125 percent of last year.

RIVER BASIN	% OF AVERAGE	% OF LAST YEAR
COLUMBIA	79	69
KOOTENAI	76	60
FLATHEAD	80	74
UPPER CLARK FORK	115	101
BITTERROOT	96	107
LOWER CLARK FORK	104	101
MISSOURI	115	126
JEFFERSON	106	99
MADISON	94	86
GALLATIN	104	92
MISSOURI MAINSTEM	116	129
SMITH-JUDITH-MUSSELSHELL	125	122
SUN-TETON-MARIAS	112	116
MILK	131	101
ST. MARY	86	111
YELLOWSTONE	113	99
UPPER YELLOWSTONE	106	90
LOWER YELLOWSTONE	113	99
STATEWIDE	105	106

Streamflow

State-wide, streamflows are forecast to be 190 percent of average. West of the Divide streamflows are forecast to be 188 percent of average and east of the Divide are forecast to be 194 percent of average. Provided near average weather conditions into the fall of 2011, streamflows and water supplies should be plentiful.

Following are streamflow forecasts for the period June 1 through July 31. THESE FORECASTS ASSUME NEAR NORMAL SPRING CONDITIONS AND DO NOT ACCOUNT FOR WELL BELOW AVERAGE (70% or less) OR WELL ABOVE AVERAGE (130% or more) SNOWMELT OR SPRING RAIN. Specific forecast probabilities are available in each individual River Basin Report.

RIVER BASIN	JUNE-JULY THIS YEAR % OF AVERAGE	JUNE-JULY LAST YEAR % OF AVERAGE
COLUMBIA	188	78
KOOTENAI	168	68
FLATHEAD	200	93
UPPER CLARK FORK	194	77
BITTERROOT	162	71
LOWER CLARK FORK	183	66
MISSOURI	191	88
JEFFERSON	183	78
MADISON	149	64
GALLATIN	156	87
MISSOURI MAINSTEM	173	79
SMITH-JUDITH-MUSSELSHELL	251	124
SUN-TETON-MARIAS	194	73
MILK	193	117
ST. MARY	155	93
YELLOWSTONE	200	88
UPPER YELLOWSTONE	170	79
LOWER YELLOWSTONE	231	101
STATE-WIDE	190	84

NOTE: The JUNE-JULY LAST YEAR % OF AVERAGE column above is what was forecast last year on June 1 and NOT what actually occurred.

Surface Water Supply Index

The Surface Water Supply Index (SWSI) is a measure of available surface water availability for the spring and summer months. Water users that rely on mountain precipitation can use the index to evaluate seasonal surface water supplies. The SWSI accounts for mountain snowpack, mountain precipitation, streamflow, reservoir storage, and soil moisture.

SWSI RATING	SURFACE WATER CONDITION
+3.0 to +4.0	Extremely Wet
+2.0 to +3.0	Moderately Wet
+1.0 to +2.0	Slightly Wet
-1.0 to +1.0	Near Average
-1.0 to -2.0	Slightly Dry
-2.0 to -3.0	Moderately Dry
-3.0 to -4.0	Extremely Dry

This Year SWSI	Last Year SWSI	Basin
+3.4	-3.2	Tobacco River
+2.4	-3.5	Kootenai Ft. Steele to Libby Dam
----	-2.2	Kootenai River below Libby Dam
+3.8	-3.0	Fisher River
+3.2	-3.5	Yaak River
+3.8	-2.3	North Fork Flathead River
+3.8	-2.4	Middle Fork Flathead River
+3.8	-0.6	South Fork Flathead River
+3.2	-2.2	Flathead River at Columbia Falls
+3.9	0.0	Swan River
+3.8	-2.3	Flathead River at Polson
----	----	Mission Valley
----	----	Little Bitterroot River
+3.5	-1.4	Clark Fork River above Milltown
+4.0	-2.7	Blackfoot River
+2.5	-2.8	Bitterroot River
+3.2	-2.4	Clark Fork River below Bitterroot River
+3.6	-2.3	Clark Fork River below Flathead River
+2.3	+0.1	Beaverhead River
+2.4	-0.7	Ruby River
+3.7	+0.2	Big Hole River
+3.0	-1.0	Boulder River (Jefferson)
+3.0	+0.7	Jefferson River
+2.8	-2.6	Madison River
+3.4	-1.0	Gallatin River
+3.3	-0.7	Missouri River above Canyon Ferry
+3.1	-1.0	Missouri River below Canyon Ferry
----	+1.7	Smith River
+3.2	-2.9	Sun River
+2.3	-0.2	Teton River
+3.3	-2.0	Birch/Dupuyer Creeks
----	+2.9	Upper Judith River
+3.4	-1.9	Marias River above Tiber
+3.9	-2.2	Marias River below Tiber
+3.8	+1.2	Musselshell River
+2.3	-0.8	Missouri River above Ft. Peck
+3.9	-1.0	Missouri River below Ft. Peck
+3.3	-1.4	St. Mary River
+3.9	+2.6	Milk River
+2.8	-1.9	Dearborn River near Craig
+3.7	-2.9	Yellowstone River above Livingston
+3.8	+1.1	Shields River
+3.6	-1.0	Boulder River (Yellowstone)
+3.8	-2.3	Stillwater River
+3.8	-0.7	Rock/Red Lodge Creeks
+3.9	-3.1	Clarks Fork River

+3.7	-2.6	Yellowstone River above Bighorn River
+4.0	-0.4	Bighorn River below Bighorn Lake
+3.8	-0.1	Little Bighorn River
+3.8	-1.6	Yellowstone River below Bighorn River
+3.9	+2.3	Tongue River
+4.0	+2.0	Powder River

Kootenai River Basin in Montana

Snowpack conditions in the Kootenai River Basin were well above average and increased 80 percent from May 1. Snow water content was 249 percent of average and 289 percent of last year. Snowpack in the Kootenai in Canada was above average. Snow water content was 146 percent of average and 163 percent of last year.



January maximum swe was established in 1997 and minimum was in 1977; February maximum swe was in 1997 and minimum swe was in 1977; March maximum swe was in 1972 and minimum swe was in 1977; April maximum swe was in 1974 and minimum swe was in 1977and 2005; May maximum swe was in 1974 and minimum swe was in 1977; and June maximum swe was in 1974 and minimum swe was in 1992. Average is for the period 1971 through 2000.

Mountain precipitation during May was 117 percent of average and 80 percent of last year. Valley precipitation during May was 100 percent of average and 97 percent of last year. Mountain and valley water year precipitation, beginning October 1, 2010, was 128 percent of average and 151 percent of last year.



Lake Koocanusa storage was 76 percent of average and 60 percent of last year.

Assuming average precipitation, June through July, streamflows are forecast to average 168 percent.

KOOTENAI RIVER BASIN in Montana
Streamflow Forecasts - June 1, 2011

Forecast Point	Forecast Period	<<= Drier = Future Conditions = Wetter =>> Chance Of Exceeding * 90% (1000AF)	70% (1000AF)	50% (1000AF)	50% (% AVG.)	30% (1000AF)	10% (1000AF)	30-Yr Avg. (1000AF)
Tobacco R nr Eureka	JUN-JUL	91	102	109	176	116	127	62
	JUN-SEP	107	120	129	170	138	151	76
Libby Reservoir Inflow (1,2)	JUN-JUL	4040	4480	4680	136	4880	5320	3440
	JUN-SEP	5010	5560	5810	131	6060	6610	4440
Fisher River nr Libby	JUN-JUL	104	116	124	182	132	144	68
	JUN-SEP	127	141	150	179	159	173	84
Yaak River nr Troy	JUN-JUL	230	255	275	191	295	320	144
	JUN-SEP	250	280	300	181	320	350	166
Kootenai R at Leonia (1,2)	JUN-JUL	5127	5720	5990	153	6260	6853	3920
	JUN-SEP	6209	6932	7260	145	7588	8311	5000

KOOTENAI RIVER BASIN in Montana
Reservoir Storage (1000 AF) - End of May

Reservoir	Usable Capacity	*** Usable Storage *** This Year	Last Year	Avg
LAKE KOOCANUSA	5748.0	2421.0	4057.0	3193.1

KOOTENAI RIVER BASIN in Montana
Watershed Snowpack Analysis - June 1, 2011

Watershed	Number of Data Sites	This Year as % of Last Yr	Average
KOOTENAY in CANADA	7	163	146
KOOTENAI MAINTSTEM	4	378	263
TOBACCO	2	190	236
FISHER	1	0	0
YAAK	3	304	274
KOOTENAI in MONTANA	9	289	249
KOOTENAI ab BONNERS FERRY	16	256	198

* 90%, 70%, 50%, 30%, and 10% chances of exceeding are the probabilities that the actual volume will exceed the volumes in the table.

The average is computed for the 1971-2000 base period.

(1) - The values listed under the 10% and 90% Chance of Exceeding are actually 5% and 95% exceedance levels.
(2) - The value is natural volume - actual volume may be affected by upstream water management.
(3) - Median value used in place of average.

Flathead River Basin

Snowpack conditions in the Flathead River Basin were well above average and increased 55 percent from May 1. Snow water content was 235 percent of average and 196 percent of last year. Snowpack in the Flathead of Canada is well above average. Snow water content was 190 percent of average and 286 percent of last year.



January maximum swe was established in 1997 and minimum was in 2001; February maximum swe was in 1997 and minimum was in 2001; March maximum swe was in 1972 and minimum was in 2005; April maximum swe was in 1972 and minimum was in 2005; May maximum swe was in 1972 and minimum was in 1992; and June maximum swe was in 1974 and minimum was in 1992. Average is for the period 1971 through 2000.

Mountain precipitation during May was 111 percent of average and 80 percent of last year. Valley precipitation during May was 86 percent of average and 77 percent of last year. Mountain and valley water year precipitation, beginning October 1, 2010, was 132 percent of average and 143 percent of last year.



The combined Camas storage was 93 percent of average; and combined Mission Valley storage was 68 percent of average; and Hungry Horse storage was 68 percent of average and 58 percent of last year; and Flathead Lake storage was 100 percent of average and 98 percent of last year.

Assuming average precipitation, June through July, streamflows are forecast to average 200 percent.

FLATHEAD RIVER BASIN
Streamflow Forecasts - June 1, 2011

Forecast Point	Forecast Period	<<= Drier = Future Conditions = Wetter =>> Chance Of Exceeding * 90% (1000AF)	70% (1000AF)	50% (1000AF)	50% (% AVG.)	30% (1000AF)	10% (1000AF)	30-Yr Avg. (1000AF)
NF Flathead R nr Columbia Falls	JUN-JUL	1330	1440	1510	182	1580	1690	830
	JUN-SEP	1540	1670	1750	175	1830	1960	1000
MF Flathead R nr West Glacier	JUN-JUL	1330	1440	1520	184	1600	1710	825
	JUN-SEP	1520	1650	1730	177	1810	1940	975
SF Flathead R nr Hungry Horse	JUN-JUL	1300	1370	1420	219	1470	1540	650
	JUN-SEP	1410	1490	1540	211	1590	1670	730
Hungry Horse Reservoir Inflow (1,2)	JUN-JUL	1830	2000	2080	217	2160	2330	960
	JUN-SEP	1990	2170	2260	207	2350	2530	1090
Flathead R at Columbia Falls (2)	JUN-JUL	4650	4980	5200	193	5420	5750	2690
	JUN-SEP	5260	5620	5870	186	6120	6480	3160
Ashley Ck nr Marion (2)	JUN-JUL	2.4	3.1	3.6	224	4.1	4.8	1.6
	JUN-SEP	2.3	3.4	4.2	382	5.0	6.1	1.1
Swan R nr Bigfork	JUN-JUL	490	520	540	177	560	590	305
	JUN-SEP	595	630	655	170	680	715	385
Flathead Lake Inflow (1,2)	JUN-JUL	5180	5730	5980	193	6230	6780	3100
	JUN-SEP	5850	6480	6770	187	7060	7690	3620
Mill Ck ab Bassoo Ck nr Niarada	JUN-JUL	2.6	2.9	3.1	246	3.3	3.6	1.3
	JUN-SEP	3.2	3.6	3.8	238	4.0	4.4	1.6
South Crow Ck nr Ronan	JUN-JUL	9.9	10.9	11.6	181	12.3	13.3	6.4
	JUN-SEP	11.7	12.9	13.7	176	14.5	15.7	7.8
Mission Ck nr St. Ignatius	JUN-JUL	25	27	28	161	29	31	17.4
	JUN-SEP	31	33	35	158	37	39	22
Sf Jocko R nr Arlee	JUN-JUL	31	33	35	212	37	39	16.5
	JUN-SEP	37	39	41	201	43	45	20
NF Jocko R bl Tabor Feeder Canal	JUN-JUL	29	31	32	216	33	35	14.8
	JUN-SEP	31	33	34	205	35	37	16.6

FLATHEAD RIVER BASIN
Reservoir Storage (1000 AF) - End of May

Reservoir	Usable Capacity	*** Usable Storage *** This Year	Last Year	Avg
CAMAS (4)	45.2	28.9	---	31.2
LOWER JOCKO LAKE	6.4	4.0	---	3.5
MISSION VALLEY (8)	0.0	0.0	---	68.7
HUNGRY HORSE	3451.0	1749.0	2999.0	2588.0
FLATHEAD LAKE	1791.0	1506.0	1531.0	1499.2

FLATHEAD RIVER BASIN
Watershed Snowpack Analysis - June 1, 2011

Watershed	Number of Data Sites	This Year as % of Last Yr	Average
NF FLATHEAD in CANADA	2	201	282
NF FLATHEAD in MONTANA	6	200	243
MIDDLE FORK FLATHEAD	4	193	187
SOUTH FORK FLATHEAD	2	182	232
STILLWATER-WHITEFISH	5	222	323
SWAN	4	169	214
MISSION VALLEY	2	152	205
LITTLE BITTERROOT-ASHLEY	0	0	0
JOCKO	5	222	254
FLATHEAD in MONTANA	21	196	235
FLATHEAD RIVER BASIN	23	197	239

* 90%, 70%, 50%, 30%, and 10% chances of exceeding are the probabilities that the actual volume will exceed the volumes in the table.

The average is computed for the 1971-2000 base period.

(1) - The values listed under the 10% and 90% Chance of Exceeding are actually 5% and 95% exceedance levels.
(2) - The value is natural volume - actual volume may be affected by upstream water management.
(3) - Median value used in place of average.

Upper Clark Fork River Basin

Snowpack conditions in the Upper Clark Fork River Basin were well above average and increased 86 percent from May 1. Snow water content was 240 percent of average and 205 percent of last year.



January maximum swe was established in 1997 and minimum swe was in 1977; February maximum was in 1972 and minimum swe was in 1977; March maximum swe was in 1972 and minimum swe was in 2005; April maximum swe was in 1972 and minimum was in 2005; May maximum swe was in 1972 and minimum swe was in 1977; and June maximum swe was in 1975 and minimum swe was in 1987. Average is for the period 1971 through 2000.

Mountain precipitation during May was 124 percent of average and 104 percent of last year. Valley precipitation during May was 104 percent of average and 95 percent of last year. Mountain and valley water year precipitation, beginning October 1, 2010, was 124 percent of average and 149 percent of last year.



East Fork Rock Creek storage was 127 percent of average and 96 percent of last year; and Nevada Creek storage was 115 percent of average and 101 percent of last year.

Assuming average precipitation, June through July, streamflows are forecast to average 194 percent.

UPPER CLARK FORK RIVER BASIN
Streamflow Forecasts - June 1, 2011

Forecast Point	Forecast Period	<<= Drier = Future Conditions = Wetter =>> Chance Of Exceeding * 90% (1000AF)	70% (1000AF)	50% (1000AF)	50% (% AVG.)	30% (1000AF)	10% (1000AF)	30-Yr Avg. (1000AF)
Little Blackfoot R nr Garrison	JUN-JUL	51	59	65	203	71	79	32
	JUN-SEP	63	73	80	205	87	97	39
Flint Ck nr Southern Cross	JUN-JUL	11.1	13.3	14.8	192	16.3	18.5	7.7
	JUN-SEP	14.9	17.9	19.9	195	22	25	10.2
Flint Ck bl Boulder Ck	JUN-JUL	45	53	59	179	65	73	33
	JUN-SEP	62	73	80	167	87	98	48
Lower Willow Ck Reservoir Inflow (2)	JUN-JUL	7.6	9.0	9.9	230	10.8	12.2	4.3
	JUN-SEP	8.9	10.4	11.5	217	12.6	14.1	5.3
MF Rock Ck nr Philipsburg	JUN-JUL	57	65	70	180	75	83	39
	JUN-SEP	66	74	80	170	86	94	47
Rock Ck nr Clinton	JUN-JUL	230	260	280	189	300	330	148
	JUN-SEP	270	305	330	180	355	390	183
Clark Fork R ab Milltown	JUN-JUL	470	545	595	189	645	720	315
	JUN-SEP	605	695	755	184	815	905	410
Nevada Ck nr Helmville	JUN-JUL	10.6	12.0	13.0	186	14.0	15.4	7.0
	JUN-SEP	12.3	13.9	15.0	177	16.1	17.7	8.5
Blackfoot R nr Bonner	JUN-JUL	670	715	745	196	775	820	380
	JUN-SEP	800	845	880	187	915	960	470
Clark Fork R ab Missoula	JUN-JUL	1170	1270	1340	193	1410	1510	695
	JUN-SEP	1430	1550	1630	185	1710	1830	880

UPPER CLARK FORK RIVER BASIN
Reservoir Storage (1000 AF) - End of May

Reservoir	Usable Capacity	*** Usable Storage *** This Year	Last Year	Avg
EAST FORK ROCK CREEK	15.6	13.3	13.8	10.5
GEORGETOWN LAKE		NO REPORT		
LOWER WILLOW CREEK		NO REPORT		
NEVADA CREEK	12.6	11.5	10.8	11.1

UPPER CLARK FORK RIVER BASIN
Watershed Snowpack Analysis - June 1, 2011

Watershed	Number of Data Sites	This Year as % of Last Yr	Average
CLARK FORK ab FLINT CREEK	5	172	237
FLINT CREEK	4	185	431
ROCK CREEK	2	209	211
CLARK FORK ab BLACKFOOT	9	195	253
BLACKFOOT	6	247	257
UPPER CLARK FORK BASIN	15	205	240

* 90%, 70%, 50%, 30%, and 10% chances of exceeding are the probabilities that the actual volume will exceed the volumes in the table.

The average is computed for the 1971-2000 base period.

(1) - The values listed under the 10% and 90% Chance of Exceeding are actually 5% and 95% exceedance levels.
(2) - The value is natural volume - actual volume may be affected by upstream water management.
(3) - Median value used in place of average.

Bitterroot River Basin

Snowpack conditions in the Bitterroot River Basin were well above average and increased 57 percent from May 1. Snow water content was 207 percent of average and 381 percent of last year.



January maximum swe was established in 1997 and minimum swe in 1977; February maximum swe was in 1972 and minimum was in 1977; March maximum swe was in 1972 and minimum swe was in 1977 and 2005; April maximum swe was in 1972 and minimum swe was in 2005; May maximum swe was in 1972 and minimum swe was in 1987; and June maximum swe was 1972 and 1974 and 1974 and minimum swe was in 1987, 1992, 1994, and 2001. Average is for the period 1971 through 2000.

Mountain precipitation during May was 120 percent of average and 96 percent of last year. Valley precipitation during May was 132 percent of average and 118 percent of last year. Mountain and valley water year precipitation, beginning October 1, 2010, was 119 percent of average and 166 percent of last year.



Painted Rocks Lake storage was 109 percent of average and 102 percent of last year and Como storage was 83 percent of average and 114 percent of last year.

Assuming near average precipitation, June through July, streamflows are forecast to average 162 percent.

BITTERROOT RIVER BASIN
Streamflow Forecasts - June 1, 2011

Forecast Point	Forecast Period	<<====== Drier ====== Future Conditions ====== Wetter =====>> Chance Of Exceeding * 90% (1000AF)	70% (1000AF)	50% (1000AF)	50% (% AVG.)	30% (1000AF)	10% (1000AF)	30-Yr Avg. (1000AF)
WF Bitterroot R nr Conner (2)	JUN-JUL	115	123	129	187	135	143	69
	JUN-SEP	126	138	146	178	154	166	82
Bitterroot R nr Darby	JUN-JUL	350	370	385	160	400	420	240
	JUN-SEP	420	445	465	158	485	510	295
Como Reservoir Inflow (2)	JUN-JUL	45	53	58	145	63	71	40
	JUN-SEP	50	58	64	146	70	78	44
Bitterroot R nr Missoula	JUN-JUL	950	1030	1080	157	1130	1210	690
	JUN-SEP	1110	1200	1260	158	1320	1410	800

BITTERROOT RIVER BASIN
Reservoir Storage (1000 AF) - End of May

Reservoir	Usable Capacity	*** Usable Storage *** This Year	Last Year	Avg
PAINTED ROCKS LAKE	31.7	33.1	32.4	30.5
COMO	34.9	25.7	22.6	30.9

BITTERROOT RIVER BASIN
Watershed Snowpack Analysis - June 1, 2011

Watershed	Number of Data Sites	This Year as % of Last Yr	Average
WEST FORK BITTERROOT	2	298	226
EAST SIDE BITTERROOT	3	299	189
WEST SIDE BITTERROOT	3	475	213
BITTERROOT RIVER BASIN	7	381	207

* 90%, 70%, 50%, 30%, and 10% chances of exceeding are the probabilities that the actual volume will exceed the volumes in the table.

The average is computed for the 1971-2000 base period.

(1) - The values listed under the 10% and 90% Chance of Exceeding are actually 5% and 95% exceedance levels.
(2) - The value is natural volume - actual volume may be affected by upstream water management.
(3) - Median value used in place of average.

Lower Clark Fork River Basin

Snowpack conditions in the Lower Clark Fork River Basin were well above average and increased 121 percent from May 1. Snow water content was 288 percent of average and 413 percent of last year.



January maximum swe was established in 1997 and minimum swe was in 1977; February maximum swe was in 1972 and minimum swe was in 1977; March maximum swe was in 1972 and minimum was in 2005; April maximum swe was in 1972 and minimum swe was in 2005; May maximum swe was in 1972 and minimum swe was in 1977; and June maximum swe was in 1974 and minimum swe was in 1977. Average is for the period 1971 through 2000.

Mountain precipitation during May was 113 percent of average and 85 percent of last year. Valley precipitation during May was 92 percent of average and 102 percent of last year. Mountain and valley water year precipitation, beginning October 1, 2010, was 126 percent of average and 167 percent of last year.



Noxon Rapids storage was 104 percent of average and 101 percent of last year.

Assuming average precipitation, June through July, streamflows are forecast to average 183 percent.

LOWER CLARK FORK RIVER BASIN
Streamflow Forecasts - June 1, 2011

Forecast Point	Forecast Period	<<= Drier = Future Conditions = Wetter =>> Chance Of Exceeding * 90% (1000AF)	70% (1000AF)	50% (1000AF)	50% (% AVG.)	30% (1000AF)	10% (1000AF)	30-Yr Avg. (1000AF)
Clark Fork R bl Missoula	JUN-JUL	2150	2310	2420	175	2530	2690	1380
	JUN-SEP	2580	2770	2890	172	3010	3200	1680
Clark Fork R at St. Regis (1)	JUN-JUL	2630	2930	3070	173	3210	3510	1770
	JUN-SEP	3170	3510	3670	170	3830	4170	2160
Clark Fork R nr Plains (1,2)	JUN-JUL	8340	9030	9350	183	9670	10400	5100
	JUN-SEP	9590	10400	10800	177	11200	12000	6090
Thompson R nr Thompson Falls	JUN-JUL	129	140	147	179	154	165	82
	JUN-SEP	161	174	183	169	192	205	108
Prospect Ck at Thompson Falls	JUN-JUL	78	81	84	205	87	90	41
	JUN-SEP	89	93	96	192	99	103	50
Clark Fork at Whitehorse Rpds (1,2)	JUN-JUL	9212	9960	10300	183	10640	11388	5620
	JUN-SEP	10604	11495	11900	176	12305	13196	6750

LOWER CLARK FORK RIVER BASIN
Reservoir Storage (1000 AF) - End of May

Reservoir	Usable Capacity	*** Usable Storage *** This Year	Last Year	Avg
NOXON RAPIDS	335.0	326.8	323.7	313.6

LOWER CLARK FORK RIVER BASIN
Watershed Snowpack Analysis - June 1, 2011

Watershed	Number of Data Sites	This Year as % of Last Yr	Average
LOWER CLARK FORK BASIN	8	413	288

* 90%, 70%, 50%, 30%, and 10% chances of exceeding are the probabilities that the actual volume will exceed the volumes in the table.

The average is computed for the 1971-2000 base period.

(1) - The values listed under the 10% and 90% Chance of Exceeding are actually 5% and 95% exceedance levels.
(2) - The value is natural volume - actual volume may be affected by upstream water management.
(3) - Median value used in place of average.

Jefferson River Basin

Snowpack conditions in the Jefferson River Basin were well above average and increased 91 percent from May 1. Snow water content was 228 percent of average and 170 percent of last year.



January maximum swe was established in 1997 and minimum swe was in 1977; February maximum swe was in 1997 and minimum was in 1977; March maximum swe was in 1972 and minimum was in 1977; April maximum swe was in 1997 and minimum was in 1977; May maximum swe was in 1975 and minimum swe was in 1977; and June maximum swe was in 1982 and minimum in 1987. Average is for the period 1971 through 2000.

Mountain precipitation during May was 151 percent of average and 107 percent of last year. Valley precipitation during May was 107 percent of average and 83 percent of last year. Mountain and valley water year precipitation, beginning October 1, 2010, was 121 percent of average and 134 percent of last year.



Lima storage was 107 percent of average and 92 percent of last year; Clark Canyon storage was 106 percent of average and 101 percent of last year; and Ruby River storage was 102 percent of average and 101 percent of last year.

Assuming average precipitation, June through July streamflows are forecast to average 183 percent.

JEFFERSON RIVER BASIN
Streamflow Forecasts - June 1, 2011

Forecast Point	Forecast Period	<<= Drier = Future Conditions = Wetter =>> Chance Of Exceeding * 90% (1000AF)	70% (1000AF)	50% (1000AF)	50% (% AVG.)	30% (1000AF)	10% (1000AF)	30-Yr Avg. (1000AF)
Lima Reservoir Inflow (2)	JUN-JUL	44	52	58	149	64	72	39
	JUN-SEP	56	67	74	157	81	92	47
Clark Canyon Reservoir Inflow (2)	JUN-JUL	44	70	88	173	106	132	51
	JUN-SEP	73	103	123	162	143	173	76
Beaverhead R at Barretts (2)	JUN-JUL	47	85	111	156	137	175	71
	JUN-SEP	70	120	153	149	186	235	103
Ruby R Reservoir Inflow (2)	JUN-JUL	56	65	71	154	77	86	46
	JUN-SEP	73	84	91	147	98	109	62
Big Hole R at Wisdom	JUN-JUL	89	112	127	235	142	165	54
	JUN-SEP	98	124	142	225	160	186	63
Big Hole R nr Melrose	JUN-JUL	565	630	675	208	720	785	325
	JUN-SEP	615	695	750	197	805	885	380
Jefferson R nr Twin Bridges (2)	JUN-JUL	635	740	815	209	880	985	390
	JUN-SEP	700	835	930	200	1020	1150	465
Boulder R nr Boulder	JUN-JUL	44	50	55	153	60	66	36
	JUN-SEP	49	58	64	149	70	79	43
Willow Ck Reservoir Inflow (2)	JUN-JUL	13.8	16.9	19.0	176	21	24	10.8
	JUN-SEP	15.9	20	23	176	26	30	13.1
Jefferson R nr Three Forks (2)	JUN-JUL	605	720	800	216	880	995	370
	JUN-SEP	720	860	955	212	1050	1190	450

JEFFERSON RIVER BASIN
Reservoir Storage (1000 AF) - End of May

Reservoir	Usable Capacity	*** Usable Storage *** This Year	Last Year	Avg
LIMA	84.0	70.3	76.2	65.8
CLARK CANYON	255.6	171.1	169.0	161.2
RUBY RIVER	38.8	38.6	38.1	37.9

JEFFERSON RIVER BASIN
Watershed Snowpack Analysis - June 1, 2011

Watershed	Number of Data Sites	This Year as % of Last Yr	Average
BEAVERHEAD	8	197	254
RUBY	5	130	201
BIGHOLE	8	177	217
BOULDER	3	173	193
JEFFERSON RIVER BASIN	19	170	228

* 90%, 70%, 50%, 30%, and 10% chances of exceeding are the probabilities that the actual volume will exceed the volumes in the table.

The average is computed for the 1971-2000 base period.

(1) - The values listed under the 10% and 90% Chance of Exceeding are actually 5% and 95% exceedance levels.
(2) - The value is natural volume - actual volume may be affected by upstream water management.
(3) - Median value used in place of average.

Gallatin River Basin

Snowpack conditions in the Gallatin River Basin were well above average and increased 101 percent from May 1. Snow water content was 244 percent of average and 187 percent of last year.



January maximum swe was established in 1997 and minimum swe was in 1966; February maximum swe was in 1997 and minimum was in 1977; March maximum swe was in 1997 and minimum was in 1977and 2005; April maximum swe was in 1997 and minimum was in 1987; May maximum swe was in 1970 and minimum swe was in 1987; and June maximum swe was in 1975 and minimum in 2001. Average is for the period 1971 through 2000.

Mountain precipitation during May was 118 percent of average and 97 percent of last year. Valley precipitation during May was 86 percent of average and 77 percent of last year. Mountain and valley water year precipitation, beginning October 1, 2010, was 120 percent of average and 128 percent of last year.



Middle Creek storage was 104 percent of average and 92 percent of last year.

Assuming average precipitation, June through July, streamflows are forecast to average 156 percent.

GALLATIN RIVER BASIN
Streamflow Forecasts - June 1, 2011

Forecast Point	Forecast Period	<<=== Drier === Future Conditions === Wetter ===>> Chance Of Exceeding * 90% (1000AF)	70% (1000AF)	50% (1000AF)	50% (% AVG.)	30% (1000AF)	10% (1000AF)	30-Yr Avg. (1000AF)
Gallatin R nr Gateway	JUN-JUL	365	400	425	149	450	485	285
	JUN-SEP	450	490	520	144	550	590	360
Hyalite Reservoir Inflow (2)	JUN-JUL	21	22	23	154	24	25	14.9
	JUN-SEP	24	26	27	150	28	30	18.0
Gallatin R at Logan	JUN-JUL	370	425	460	164	495	550	280
	JUN-SEP	455	525	570	158	615	685	360

GALLATIN RIVER BASIN
Reservoir Storage (1000 AF) - End of May

Reservoir	Usable Capacity	*** Usable Storage *** This Year	Last Year	Avg
MIDDLE CREEK	10.2	7.9	8.6	7.6

GALLATIN RIVER BASIN
Watershed Snowpack Analysis - June 1, 2011

Watershed	Number of Data Sites	This Year as % of Last Yr	Average
UPPER GALLATIN	3	189	197
HYALITE	2	175	224
BRIDGER	2	199	741
GALLATIN RIVER BASIN	7	187	244

* 90%, 70%, 50%, 30%, and 10% chances of exceeding are the probabilities that the actual volume will exceed the volumes in the table.

The average is computed for the 1971-2000 base period.

(1) - The values listed under the 10% and 90% Chance of Exceeding are actually 5% and 95% exceedance levels.
(2) - The value is natural volume - actual volume may be affected by upstream water management.
(3) - Median value used in place of average.

Missouri Mainstem River Basin

Snowpack conditions in the Headwaters Missouri Mainstem River Basin were well above average and increased 143 percent from May 1. Snow water content was 300 percent of average and 207 percent of last year.



January maximum swe was established in 1997 and minimum swe in 1977; February maximum swe was in 1972 and minimum swe was in 1977 and 1988; March maximum swe in 1972 and minimum swe was in 1977; April maximum swe was in 1972 and minimum swe was in 1961, 1966 and 2005; May maximum swe was in 1975 and minimum swe was in 1977; and June maximum swe was in 1975 and minimum swe was in 1992. Average is for the period 1971 through 2000.

Mountain precipitation during May was 134 percent of average and 104 percent of last year. Valley precipitation during May was 230 percent of average and 134 percent of last year. Mountain and valley water year precipitation, beginning October 1, 2010, was 145 percent of average and 125 percent of last year.



Canyon Ferry Lake storage was 85 percent of average and 83 percent of last year; Helena Valley storage was 114 percent of average and 97 percent of last year; Lake Helena storage was 76 percent of average and 99 percent of last year; Hauser & Helena storage was 118 percent of average and 100 percent of last year; Holter Lake storage was 103 percent of average and 101 percent of last year; and Fort Peck Lake storage was 119 percent of average and 135 percent of last year.

Assuming average precipitation, June through July, streamflows are forecast to average 173 percent.

MISSOURI MAINSTEM RIVER BASIN
Streamflow Forecasts - June 1, 2011

Forecast Point	Forecast Period	<<= Drier = Future Conditions = Wetter =>> Chance Of Exceeding * 90% (1000AF)	70% (1000AF)	50% (1000AF)	50% (% AVG.)	30% (1000AF)	10% (1000AF)	30-Yr Avg. (1000AF)
Missouri R at Toston (2)	JUN-JUL	1520	1740	1890	173	2040	2260	1090
	JUN-SEP	1940	2250	2450	173	2650	2960	1420
Dearborn R nr Craig	JUN-JUL	83	97	107	170	117	131	63
	JUN-SEP	83	99	110	164	121	137	67
Missouri R at Fort Benton (2)	JUN-JUL	2410	2560	2670	169	2780	2930	1580
	JUN-SEP	3300	3520	3670	169	3820	4030	2170
Missouri R nr Virgelle (2)	JUN-JUL	2530	2890	3130	172	3370	3730	1820
	JUN-SEP	3410	3870	4190	172	4510	4980	2430
Missouri R nr Landusky (2)	JUN-JUL	2710	3090	3350	172	3610	3990	1950
	JUN-SEP	3610	4130	4480	172	4830	5340	2610
Missouri R bl Fort Peck Dam (2)	JUN-JUL	2790	3150	3390	173	3630	3990	1960
	JUN-SEP	3620	4080	4400	173	4720	5180	2550
Lake Sakakawea Inflow (2)	JUN-JUL	8700	9830	10600	183	11400	12500	5800
	JUN-SEP	11200	12200	13200	182	14200	15200	7240

MISSOURI MAINSTEM RIVER BASIN
Reservoir Storage (1000 AF) - End of May

Reservoir	Usable Capacity	*** Usable Storage *** This Year	Last Year	Avg
CANYON FERRY LAKE	2043.0	1403.0	1686.0	1656.7
HELENA VALLEY	9.2	8.8	9.1	7.7
LAKE HELENA	12.7	10.9	11.0	14.4
HAUSER & HELENA	74.6	74.0	74.3	62.6
HOLTER LAKE	81.9	81.8	81.3	79.8
FORT PECK LAKE	18910.0	18426.0	13660.0	15495.0

MISSOURI MAINSTEM RIVER BASIN
Watershed Snowpack Analysis - June 1, 2011

Watershed	Number of Data Sites	This Year as % of Last Yr	Average
HEADWATERS MAINSTEM	5	207	300
SMITH-JUDITH-MUSSELSHELL	9	152	386
SUN-TETON-MARIAS	6	326	273
MAINSTEM ab FT PECK RES	19	204	319
MILK RIVER BASIN	3	0	0
MISSOURI MAINSTEM BASIN	21	195	328

* 90%, 70%, 50%, 30%, and 10% chances of exceeding are the probabilities that the actual volume will exceed the volumes in the table.

The average is computed for the 1971-2000 base period.

(1) - The values listed under the 10% and 90% Chance of Exceeding are actually 5% and 95% exceedance levels.
(2) - The value is natural volume - actual volume may be affected by upstream water management.
(3) - Median value used in place of average.

Smith-Judith-Musselshell River Basins

Snowpack conditions in the Smith-Judith-Musselshell River Basins were well above average and increased 198 percent from May 1. Snow water content was 386 percent of average and 152 percent of last year. Snow water content in the Smith River Basin was 173 percent of average and 173 percent of last year; the Judith River Basin was 197 percent of average and 146 percent of last year; and the Musselshell Basin River was 177 percent of average and 235 percent of last year.



January maximum swe was established in 1997 and minimum swe in 1988; February maximum swe was in 1978 and minimum swe was in 1987; March maximum swe was in 1978 and minimum swe was in 1987 and 2005; April maximum swe was in 1970 and minimum swe was in 1992; and May maximum swe was in 1970 and minimum swe was in 1987; and June maximum swe was in 1967 and minimum swe was in 1987, 1992, 1994, and 2001. Average is for the period 1971 through 2000.

Mountain and valley precipitation during May in the Smith-Belts was 160 percent of average and 119 percent of last year; in the Judith was 246 percent of average and 186 percent of last year; and in the Musselshell was 276 percent of average and 277 percent of last year. Mountain and valley water year precipitation for the greater basin, beginning October 1, 2010, was 153 percent of average and 131 percent of last year.



Smith River storage was 114 percent of average and 102 percent of last year; Ackley storage was 116 percent of average and 89 percent of last year; Bair storage was 127 percent of average and 123 percent of last year; Martinsdale storage was 109 percent of average and 101 percent of last year; and Deadman's Basin was 132 percent of average and 139 percent of last year.

Assuming average precipitation, June through July, streamflows are forecast to average 251 percent.

SMITH-JUDITH-MUSSELSHELL RIVER BASINS
Streamflow Forecasts - June 1, 2011

Forecast Point	Forecast Period	<<====== Drier ====== Future Conditions ======= Wetter =====>> Chance Of Exceeding * 90% (1000AF)	70% (1000AF)	50% (1000AF)	50% (% AVG.)	30% (1000AF)	10% (1000AF)	30-Yr Avg. (1000AF)
Sheep Ck nr White Sulphur Springs	JUN-JUL	15.6	17.8	19.3	212	21	23	9.1
	JUN-SEP	19.0	22	24	198	26	29	12.1
Smith R bl Eagle Ck (2)	JUN-JUL	126	146	160	213	174	194	75
	JUN-SEP	140	168	186	204	205	230	91
NF Musselshell R nr Delpine	JUN-JUL	3.8	5.4	6.6	287	7.8	9.4	2.3
	JUN-SEP	5.9	7.3	8.3	259	9.3	10.7	3.2
SF Musselshell R ab Martinsdale	JUN-JUL	27	38	46	184	54	65	25
	JUN-SEP	30	42	50	172	58	70	29
Musselshell R at Harlowton (2)	JUN-JUL	65	91	109	295	127	153	37
	JUN-SEP	73	99	117	285	135	161	41
Musselshell R nr Roundup (2)	JUN-JUL	106	131	148	315	165	190	47
	JUN-SEP	117	142	159	318	176	200	50

SMITH-JUDITH-MUSSELSHELL RIVER BASINS
Reservoir Storage (1000 AF) - End of May

Reservoir	Usable Capacity	*** Usable Storage *** This Year	Last Year	Avg
SMITH RIVER	10.6	11.6	11.4	10.2
ACKLEY LAKE	7.0	5.7	6.4	4.9
BAIR	7.0	7.5	6.1	5.9
MARTINSDALE	23.1	19.9	19.8	18.2
DEADMAN'S BASIN	72.2	76.2	55.0	57.6

SMITH-JUDITH-MUSSELSHELL RIVER BASINS
Watershed Snowpack Analysis - June 1, 2011

Watershed	Number of Data Sites	This Year as % of Last Yr	This Year as % of Average
SMITH	6	163	333
HIGHWOOD	0	0	0
JUDITH	4	151	383
MUSSELSHELL	2	224	432
SMITH-JUDITH-MUSSELSHELL	9	152	386

* 90%, 70%, 50%, 30%, and 10% chances of exceeding are the probabilities that the actual volume will exceed the volumes in the table.

The average is computed for the 1971-2000 base period.

(1) - The values listed under the 10% and 90% Chance of Exceeding are actually 5% and 95% exceedance levels.
(2) - The value is natural volume - actual volume may be affected by upstream water management.
(3) - Median value used in place of average.

Sun-Teton-Marias River Basins

Snowpack conditions in the Sun-Teton-Marias River Basins were well above average and increased 107 percent from May 1. Snow water content was 273 percent of average and 326 percent of last year. Snow water content in the Sun River Basin was 368 percent of average and 471 percent of last year; the Teton River Basin was 384 percent of average and 491 percent of last year; and the Marias River Basin was 225 percent of average and 262 percent of last year.



January maximum swe was established in 1997 and minimum swe was in 1988; February maximum swe was in 1972 and minimum swe was in 2001; March maximum swe was in 1972 and minimum swe was in 2005; April maximum swe was in 1972 and minimum swe was in 2005; May maximum swe was in 1972 and minimum swe was in 1977; and June maximum was in 1972 and minimum swe was in 1987, 1992, and 1998. Average is for the period 1971 through 2000.

Mountain and valley precipitation during May in the Sun was 167 percent of average and 135 percent of last year; in the Teton was 136 percent of average and 115 percent of last year; and in the Marias was 129 percent of average and 102 percent of last year. Mountain and valley water year precipitation, beginning October 1, 2010, was 124 percent of average and 144 percent of last year.



Gibson storage was 84 percent of average and 87 percent of last year; Pishkun storage was 102 percent of average and 99 percent of last year; Willow Creek storage was 113 percent of average and 106 percent of last year; Swift storage was 60 percent of average and 68 percent of last year; Lake Frances storage was 120 percent of average and 155 percent of last year; and Lake Elwell (Tiber) storage was 118 percent of average and 117 percent of last year.

Assuming average precipitation, June through July, streamflows are forecast to average 194 percent.

SUN-TETON-MARIAS RIVER BASINS
Streamflow Forecasts - June 1, 2011

Forecast Point	Forecast Period	<<====== Drier ====== Future Conditions ====== Wetter ======>> Chance Of Exceeding * 90% (1000AF)	70% (1000AF)	50% (1000AF)	50% (% AVG.)	30% (1000AF)	10% (1000AF)	30-Yr Avg. (1000AF)
Gibson Reservoir Inflow (2)	JUN-JUL	385	420	440	173	460	495	255
	JUN-SEP	465	500	525	172	550	585	305
Two Medicine R nr Browning (2)	JUN-JUL	128	142	151	159	160	174	95
	JUN-SEP	143	158	169	158	180	195	107
Badger Ck nr Browning	JUN-JUL	60	70	77	175	84	94	44
	JUN-SEP	71	83	91	172	99	111	53
Swift Reservoir Inflow (2)	JUN-JUL	55	61	65	186	69	75	35
	JUN-SEP	69	77	82	175	87	95	47
Dupuyer Ck nr Valier	JUN-JUL	10.0	14.4	17.5	269	21	25	6.5
	JUN-SEP	13.0	18.4	22	265	26	31	8.3
Cut Bank Ck nr Browning	JUN-JUL	57	65	70	167	75	83	42
	JUN-SEP	66	74	80	163	86	94	49
Marias R nr Shelby (2)	JUN-JUL	305	370	415	227	460	525	183
	JUN-SEP	315	390	440	210	490	565	210
Teton R nr Dutton	JUN-JUL	21	39	52	193	65	83	27
	JUN-SEP	32	54	69	197	84	106	35

SUN-TETON-MARIAS RIVER BASINS
Reservoir Storage (1000 AF) - End of May

Reservoir	Usable Capacity	*** Usable Storage *** This Year	Last Year	Avg
GIBSON	99.1	74.6	85.4	89.2
PISHKUN	32.0	30.8	31.1	30.3
WILLOW CREEK	32.2	31.5	29.7	28.0
LOWER TWO MEDICINE LAKE	11.9	7.9	4.9	12.3
FOUR HORNS LAKE	19.2	11.8	4.1	12.7
SWIFT	30.0	14.0	20.7	23.5
LAKE FRANCES	112.0	100.8	65.1	83.9
LAKE ELWELL (TIBER)	1347.0	868.8	743.7	737.7

SUN-TETON-MARIAS RIVER BASINS
Watershed Snowpack Analysis - June 1, 2011

Watershed	Number of Data Sites	This Year as % of Last Yr	Average
SUN	2	471	368
TETON	3	491	384
MARIAS	3	262	225
SUN-TETON-MARIAS	6	326	273

* 90%, 70%, 50%, 30%, and 10% chances of exceeding are the probabilities that the actual volume will exceed the volumes in the table.

The average is computed for the 1971-2000 base period.

(1) - The values listed under the 10% and 90% Chance of Exceeding are actually 5% and 95% exceedance levels.
(2) - The value is natural volume - actual volume may be affected by upstream water management.
(3) - Median value used in place of average.

St. Mary and Milk River Basins

Snowpack in the Saint Mary River Basin was well above average and increased 22 percent from May 1. Snow water content was 167 percent of average and 161 percent of last year. The Milk River Basin (Bearpaw Mountains) snowpack is fully melted.





Bearpaw - January maximum swe was established in 1978 and minimum swe was in 1981; February maximum swe was 1978 and minimum was in 1973; March maximum swe was 1978 and minimum swe was 2005; April maximum swe was in 1985 and minimum swe was in 1973, 1983, and 1986; May maximum swe was 1975 and the minimum, 0.0, has occurred in several years. Average is for the period 1971 through 2000.

St. Mary - January maximum swe was established in 1997 and minimum swe was in 1988; February maximum swe was in 1972 and minimum swe was in 2001; March maximum swe was in 1972 and minimum swe was in 2005; April maximum swe was in 1967 and 1972 and minimum swe was in 2005; May maximum swe was in 19972 and minimum swe was in 1977; and June maximum swe was in 1972 and minimum swe was 1992. Average is for the period 1971 through 2000.

Mountain and valley precipitation in the St. Mary River Basin during May was 104 percent of average and 89 percent of last year; and in the Milk River Basin during May was 173 percent of average and 100 percent of last year. Mountain and valley water year precipitation for both basins, beginning October 1, 2010, was 132 percent of average and 114 percent of last year.





Lake Sherburne storage was 86 percent of average and 111 percent of last year; Fresno storage was 124 percent of average and 100 percent of last year; Beaver Creek storage was not available; and Nelson storage was 145 percent of average and 101 percent of last year.

Assuming average precipitation, June through July, streamflows in the St. Mary are forecast to average 155 percent and the Milk to average 193 percent.

ST. MARY and MILK RIVER BASINS
Streamflow Forecasts - June 1, 2011

Forecast Point	Forecast Period	<<====== Drier ====== Future Conditions ====== Wetter ======>> Chance Of Exceeding * 90% (1000AF)	70% (1000AF)	50% (1000AF)	50% (% AVG.)	30% (1000AF)	10% (1000AF)	30-Yr Avg. (1000AF)
Lake Sherburne Inflow (2)	JUN-JUL	79	87	92	151	97	105	61
	JUN-SEP	97	106	112	144	118	127	78
St. Mary R nr Babb (2)	JUN-JUL	320	350	370	148	390	420	250
	JUN-SEP	385	420	445	141	470	505	315
St. Mary R at Int'l Boundary (2)	JUN-JUL	370	410	440	166	470	510	265
	JUN-SEP	450	500	530	154	560	610	345
Milk R at Western Crossing (3)	JUN-JUL	9.5	14.0	17.1	180	20	25	9.5
	JUN-SEP	10.0	15.9	20	177	24	30	11.3
Milk R at Eastern Crossing (2,3)	JUN-JUL	3.3	23	37	190	51	71	19.5
	JUN-SEP	16.4	38	53	183	68	90	29
Beaver Ck nr Havre	JUN-JUL	3.3	4.9	8.3	208	11.7	16.7	4.0

ST. MARY and MILK RIVER BASINS
Reservoir Storage (1000 AF) - End of May

Reservoir	Usable Capacity	*** Usable Storage *** This Year	Last Year	Avg
LAKE SHERBURNE	64.3	29.7	26.8	34.6
FRESNO	127.0	103.7	103.3	83.5
BEAVER CREEK		NO REPORT		
NELSON	66.8	58.9	58.1	40.6

ST. MARY and MILK RIVER BASINS
Watershed Snowpack Analysis - June 1, 2011

Watershed	Number of Data Sites	This Year as % of Last Yr	Average
ST. MARY	2	161	170
BEARPAW MOUNTAINS	3	0	0
CYPRESS HILLS, CANADA	0	0	0
MILK RIVER BASIN	3	0	0
ST. MARY & MILK BASINS	5	161	167

* 90%, 70%, 50%, 30%, and 10% chances of exceeding are the probabilities that the actual volume will exceed the volumes in the table.

The average is computed for the 1971-2000 base period.

(1) - The values listed under the 10% and 90% Chance of Exceeding are actually 5% and 95% exceedance levels.
(2) - The value is natural volume - actual volume may be affected by upstream water management.
(3) - Median value used in place of average.

Upper Yellowstone River Basin

Snowpack conditions in the Upper Yellowstone River Basin were well above average and increased 71 percent from May 1. Snow water content was 222 percent of average and 217 percent of last year.



January maximum swe was established in 1997 and minimum swe was in 1988; February maximum swe was in 1997 and minimum swe was in 2001; March maximum swe was in 1997 and minimum swe was in 2001; April maximum swe was in 1997 and minimum swe was in 2001; May maximum swe was in 1997 and minimum swe was in 1987; and June maximum swe was 1996 and minimum swe was in 2001. Average is for the period 1971 through 2000.

Mountain precipitation during May was 157 percent of average and 135 percent of last year. Valley precipitation during May was 256 percent of average and 292 percent of last year. Mountain and valley water year precipitation, beginning October 1, 2010, was 134 percent of average and 162 percent of last year.



Mystic Lake storage was 7 percent of average and 10 percent of last year and Cooney storage was 132 percent of average and 101 percent of last year.

Assuming average precipitation, June through July, streamflows are forecast to average 170 percent.

UPPER YELLOWSTONE RIVER BASIN
Streamflow Forecasts - June 1, 2011

Forecast Point	Forecast Period	<<= Drier = Future Conditions = Wetter =>> Chance Of Exceeding * 90% (1000AF)	70% (1000AF)	50% (1000AF)	50% (% AVG.)	30% (1000AF)	10% (1000AF)	30-Yr Avg. (1000AF)
Yellowstone R at Yellowstone Lake	JUN-JUL	730	775	805	166	835	880	485
	JUN-SEP	990	1060	1100	158	1140	1210	695
Yellowstone R at Corwin Springs	JUN-JUL	1670	1800	1880	165	1960	2090	1140
	JUN-SEP	2060	2220	2330	160	2440	2600	1460
Yellowstone R at Livingston	JUN-JUL	1900	2050	2150	164	2250	2400	1310
	JUN-SEP	2330	2530	2660	157	2790	2990	1700
Shields R nr Livingston	JUN-JUL	104	134	155	225	176	205	69
	JUN-SEP	119	155	180	209	205	240	86
Boulder R at Big Timber	JUN-JUL	320	345	365	178	385	410	205
	JUN-SEP	350	380	405	176	430	460	230
West Rosebud Ck nr Roscoe (2)	JUN-JUL	63	68	71	151	74	79	47
	JUN-SEP	85	92	96	150	100	107	64
Stillwater R nr Absarokee (2)	JUN-JUL	500	540	565	155	590	630	365
	JUN-SEP	605	660	695	151	730	785	460
Clarks Fk Yellowstone R nr Belfry	JUN-JUL	615	655	680	174	705	745	390
	JUN-SEP	690	740	775	174	810	860	445
Cooney Reservoir Inflow (2) (2)	JUN-JUL	24	30	34	148	38	44	23
Cooney Reservoir Inflow (2)	JUN-SEP	36	44	49	144	54	62	34
Yellowstone R at Billings (2)	JUN-JUL	3430	3800	4050	170	4300	4670	2380
	JUN-SEP	4660	4770	5090	170	5410	5520	2990

UPPER YELLOWSTONE RIVER BASIN
Reservoir Storage (1000 AF) - End of May

Reservoir	Usable Capacity	*** Usable Storage *** This Year	Last Year	Avg
MYSTIC LAKE	21.0	0.4	4.0	5.7
COONEY	27.4	28.3	27.9	21.5

UPPER YELLOWSTONE RIVER BASIN
Watershed Snowpack Analysis - June 1, 2011

Watershed	Number of Data Sites	This Year as % of Last Yr	Average
YELLOWSTONE ab LIVINGSTON	11	239	223
SHIELDS	4	179	374
BOULDER-STILLWATER	3	202	211
RED LODGE-ROCK CREEK	2	145	229
CLARK'S FORK	7	221	184
UPPER YELLOWSTONE BASIN	23	217	222

* 90%, 70%, 50%, 30%, and 10% chances of exceeding are the probabilities that the actual volume will exceed the volumes in the table.

The average is computed for the 1971-2000 base period.

(1) - The values listed under the 10% and 90% Chance of Exceeding are actually 5% and 95% exceedance levels.
(2) - The value is natural volume - actual volume may be affected by upstream water management.
(3) - Median value used in place of average.

Lower Yellowstone River Basin

Snowpack conditions in the Lower Yellowstone River Basin were well above average and increased 166 percent from May 1. Snow water content was 305 percent of average and 227 percent of last year.



January maximum swe was established in 1997 and minimum swe was in 2000; February maximum swe was in 1997 and minimum swe was in 2001; March maximum swe was in 1986 and minimum swe was in 2001; April maximum swe was in 1986 and minimum swe was in 2001; May maximum swe was in 1997 and minimum swe was in 1981; and June maximum swe was in 1995 and minimum swe was in 2001. Average is for the period 1971 through 2000.

Mountain and valley precipitation during May was 261 percent of average and 149 percent of last year. Mountain and valley water year precipitation, beginning October 1, 2010, was 146 percent of average and 149 percent of last year.



Bighorn Lake storage was 110 percent of average and 99 percent of last year and Tongue River storage was 170 percent of average and 100 percent of last year.

Assuming average precipitation, June through July, streamflows are forecast to average 231 percent.

LOWER YELLOWSTONE RIVER BASIN
Streamflow Forecasts - June 1, 2011

Forecast Point	Forecast Period	<<====== Drier ====== Future Conditions ======= Wetter =====>> Chance Of Exceeding * 90% (1000AF)	70% (1000AF)	50% (1000AF)	50% (% AVG.)	30% (1000AF)	10% (1000AF)	30-Yr Avg. (1000AF)
Bighorn R nr St. Xavier (2)	JUN-JUL	1840	2030	2240	206	2270	2460	1090
	JUN-SEP	2140	2380	2630	212	2700	2940	1240
Little Bighorn R nr Hardin	JUN-JUL	128	145	157	234	169	186	67
	JUN-SEP	148	168	182	219	196	215	83
Tongue R nr Dayton (2)	JUN-JUL	109	119	125	216	131	141	58
	JUN-SEP	126	137	145	204	153	164	71
Big Goose Ck nr Sheridan	JUN-JUL	74	79	83	237	87	92	35
	JUN-SEP	82	88	92	209	96	102	44
Little Goose Ck nr Bighorn	JUN-JUL	42	45	47	224	49	52	21
	JUN-SEP	52	55	58	200	61	64	29
Tongue River Reservoir Inflow (2)	JUN-JUL	275	300	320	254	340	365	126
	JUN-SEP	305	340	360	235	380	415	153
Yellowstone R at Miles City (2)	JUN-JUL	5360	5920	6300	175	6680	7240	3600
	JUN-SEP	7250	7310	7780	175	8250	8310	4450
Powder R at Moorhead	JUN-JUL	240	275	295	281	315	350	105
	JUN-SEP	275	315	340	266	365	405	128
Powder R nr Locate	JUN-JUL	265	305	335	289	365	405	116
	JUN-SEP	300	355	390	277	425	480	141
Yellowstone R nr Sidney (2)	JUN-JUL	6010	6580	6970	191	7360	7930	3650
	JUN-SEP	7800	8030	8500	191	8970	9200	4460

LOWER YELLOWSTONE RIVER BASIN
Reservoir Storage (1000 AF) - End of May

Reservoir	Usable Capacity	*** Usable Storage *** This Year	Last Year	Avg
BIGHORN LAKE	1356.0	956.3	962.3	867.1
TONGUE RIVER	79.1	81.6	81.8	48.0

LOWER YELLOWSTONE RIVER BASIN
Watershed Snowpack Analysis - June 1, 2011

Watershed	Number of Data Sites	This Year as % of Last Yr	Average
WIND RIVER (Wyoming)	12	198	310
SHOSHONE RIVER (Wyoming)	6	231	201
BIGHORN RIVER (Wyoming)	16	257	269
LITTLE BIGHORN (Wyoming)	2	219	293
TONGUE RIVER (Wyoming)	7	239	452
POWDER RIVER (Wyoming)	6	338	616
LOWER YELLOWSTONE BASIN (	31	227	305

* 90%, 70%, 50%, 30%, and 10% chances of exceeding are the probabilities that the actual volume will exceed the volumes in the table.

The average is computed for the 1971-2000 base period.

(1) - The values listed under the 10% and 90% Chance of Exceeding are actually 5% and 95% exceedance levels.
(2) - The value is natural volume - actual volume may be affected by upstream water management.
(3) - Median value used in place of average.

Issued by

Dave White
Chief
Natural Resources Conservation Service
U.S. Department of Agriculture

Released by

Joyce Swartzendruber
State Conservationist
Natural Resources Conservation Service
Bozeman, Montana



Federal Building, Room 443
10 E. Babcock
Bozeman, MT 59715

Montana
Water Supply Outlook Report

Natural Resources Conservation Service
Bozeman, MT